

Discussion of
In pursuit of balance
David McKenzie, Miriam Bruhn

Markus Frölich

Randomized controlled trials

Randomized experiments now very popular

But implementing random experiments is not taught in econometrics

Basic idea: 1) Randomly allocate people to control versus treatment
2) and compare **average outcome** of treated and control

But inefficient (particularly with small samples)

Ex post balance: e.g. by regression methods on X

Ex ante balance: stratification, pair-matching, re-randomization

→ This paper - summarizes some results for design in **clinical trials**
- examines balance and inference with 4 typical datasets

Randomization methods:

- Stratification (also on Y_{t-1} !)
- Pair-wise matching (Mahalanobis distance)
- Re-randomization methods

→ Overall **pair-wise matching** seems to perform best

→ Include rather many variables in X

1) Optimal greedy algorithm takes several days with $n=300$ in Stata

→ Develop faster code and share it with researchers

2) Pair-wise matching **combined** with re-randomization is missing

Overall **pair-wise matching** seems to perform best

- in achieving balance and also with respect to power of test

For inference important to include randomization variable in regression

When pair-wise matching, include a dummy for each pair

$$Y_i = \alpha + \beta D_i + \xi_1 \text{Pair}_1 + \xi_2 \text{Pair}_2 + \xi_3 \text{Pair}_3 \dots$$

But: How to proceed in non-linear model (**incidental parameters** problem)

- Compare **randomization based inference** to **model based inference**

Implementation of randomized controlled trials

Hence: Implementation of experiments not so trivial (with small samples)

Randomization methods matter and should be accounted for in inference

Yet in practice more problems may occur:

- Non-compliance with protocol (local average treatment effect)
- Changing treatment dose over time
- Combinations of treatment over time

- Missing baseline data
- Attrition, non-response and loss to follow up (particularly long-term)

Not everything under control of experimenter

Build in multiple IV to control for these problems

Learn more from biometrics/biostats literature on longitudinal trials

But: biostats/clinical trials like ML and not IV

- Need to learn more about experiments in imperfect conditions
- Paper McKenzie & Bruhn right direction
- How to do inference in IV setup with heterogenous Treatment Effects